KEY TO THE COMMON FERNS OF MARYLAND

Prepared by Dwight Johnson, September 2014

I. Broadest at Base (Broadly Triangular in Shape)	
Once cut with wavy pinnae	Sensitive Fern
Twice cut with wiry stipe	Broad Beech Fern
• Thrice cut	
i. Large, coarse; usually more than one plant in area; not lacy	Bracken Fern
ii. Small and growing singly or in a small group; delicate and lacy	
A. Fertile stalk arising from center of plant; on lowest pinna,	
the pinnule closest to the stipe is shorter than the second pinnule	Rattlesnake Fern
B. Fertile stalk arising from base of plant; on lowest pinna,	
the pinnule closest to the stipe is longer than the second pinnule	Cut Leaf Grape Fern
II. Fully Tapering to the Base	
• Once cut	
Small-medium size; very narrow frond; pinnae oblong with "boot" similar	
to Christmas fern; stipe smooth and dark	Ebony Spleenwort
• Twice cut	
i. Large and growing in a definite crown; fertile frond (if present) in center	er,
found more often in gardens	
ii. Medium size, and growing zig-zag over an area, not in crowns	New York Fern
III. Semi-tapering to the base (Lanced Shaped)	
• Once cut	
i. Pinnae two inches or less	
A. Boot-shaped pinnae attached by a short stalk;	Christmas Fern
B. Pinnae with no stalk; usually growing on top of rocks	Common Polypody
ii. Pinnae over two inches long, wavy (resembles Sensitive Fern),	
found in moist areas east of I-95	Netted Chain Fern
• Twice Cut	
i. Stipe with brown scales; sori on edge of pinnule	Marginal Wood Fern
ii. Stipe smooth or slightly hairy	
A. Large size; woolly tufts at bases (axes) of pinnae	Cinnamon Fern
B. No tufts at axes	
1. Large size; some pinnae apparently missing from	
middle of fertile fronds (if present); veins in pinnule forked	
2. Sori and pinnule veination in herringbone pattern; 2 lowest pinn	
pointing outward in a V shape	
3. Growing zigzag in patches near wet areas; common east of I-95	
4. Large and shrubby; leaf-like pinnules with short stems	
• Thrice Cut	
i. Stipe with brown scales	
A. On lowest pinna, the pinnule closest to the stipe is clearly longer	Spinulose Wood Form
than the second pinnule B. On lowest pinna, the pinnule closest to the stipe is about the	Spinulose wood Fern
the same length or shorter than the second pinnule	Intermediate Wood Forn
ii. Stipe smooth	Three mediate wood Fern
A. Growing in a clump; sori slightly curved and in herringbone pattern	1:
rachis smooth, not hairy	
B. Growing zig-zag over area; hairy rachis; sori small and round	
C. Pinnules lobed (leaf-like); stipe long and slender	
iii. Stipe and rachis hairy, pinnules lobed; lower pinnules rounded,.	
small fern found growing in rocks	Blunt-lobed Woodsia
- · · · · · · · · · · · · · · · · · · ·	
IV. Unique Shaped Fern	
• Circular-shaped fronds with pinnae on outer rim of stalk; dark wiry stipe	Maidenhair Fern